



wwPDB X-ray Structure Validation Summary Report

Oct 13, 2024 – 01:39 pm BST

PDB ID : 1QKC
Title : ESCHERICHIA COLI FERRIC HYDROXAMATE UPTAKE RECEPTOR (FHUA) IN COMPLEX DELTA TWO-ALBOMYCIN
Authors : Ferguson, A.D.; Braun, V.; Fiedler, H.-P.; Coulton, J.W.; Diederichs, K.; Welte, W.
Deposited on : 1999-07-18
Resolution : 3.10 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the  symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references](#) ) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.4, CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 3.0
buster-report : 1.1.7 (2018)
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4 : 9.0.003 (Gargrove)
Density-Fitness : 1.0.11
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.39


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	180529	1454 (3.10-3.10)
Ramachandran outliers	177936	1391 (3.10-3.10)
Sidechain outliers	177891	1391 (3.10-3.10)
RSRZ outliers	164620	1351 (3.10-3.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	725	
2	B	11	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	GMH	B	10	X	-	-	-
2	GCN	B	2	X	-	-	-
2	GMH	B	5	X	-	-	-
3	FTT	A	1006	X	-	-	-
5	PO4	A	1014	-	-	X	-
7	ALB	A	1022[A]	X	-	-	-
7	ALB	A	1022[B]	X	-	-	-

2 Entry composition i

There are 8 unique types of molecules in this entry. The entry contains 6195 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called FERRIC HYDROXAMATE RECEPTOR.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	707	5523	3475	944	1090	14	0	0	0

There are 11 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	406	SER	-	insertion	UNP P06971
A	407	SER	-	insertion	UNP P06971
A	408	HIS	-	insertion	UNP P06971
A	409	HIS	-	insertion	UNP P06971
A	410	HIS	-	insertion	UNP P06971
A	411	HIS	-	insertion	UNP P06971
A	412	HIS	-	insertion	UNP P06971
A	413	HIS	-	insertion	UNP P06971
A	414	GLY	-	insertion	UNP P06971
A	415	SER	-	insertion	UNP P06971
A	416	SER	-	insertion	UNP P06971

- Molecule 2 is an oligosaccharide called alpha-D-glucopyranose-(1-2)-alpha-D-glucopyranose-(1-3)-[alpha-D-galactopyranose-(1-6)]alpha-D-glucopyranose-(1-3)-[L-glycero-alpha-D-manno-heptopyranose-(1-7)]L-glycero-alpha-D-manno-heptopyranose-(1-3)-L-glycero-alpha-D-manno-heptopyranose-(1-5)-[3-deoxy-alpha-D-manno-oct-2-ulopyranosonic acid-(2-4)]3-deoxy-alpha-D-manno-oct-2-ulopyranosonic acid-(2-6)-2-amino-2,3-dideoxy-alpha-D-glucopyranose-(1-6)-2-amino-2-deoxy-alpha-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
2	B	11	134	73	2	59	0	0	0

- Molecule 3 is 3-HYDROXY-TETRADECANOIC ACID (three-letter code: FTT) (formula: $C_{14}H_{28}O_3$).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C O 16 14 2	0	0
3	A	1	Total C O 7 4 3	0	0
3	A	1	Total C O 16 14 2	0	0
3	A	1	Total C O 13 12 1	0	0
3	A	1	Total C O 17 14 3	0	0
3	A	1	Total C O 15 14 1	0	0

- Molecule 4 is DIPHOSPHATE (three-letter code: DPO) (formula: O_7P_2).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	O	P	0	0
			8	6	2		
4	A	1	Total	O	P	0	0
			8	6	2		

- Molecule 5 is PHOSPHATE ION (three-letter code: PO4) (formula: O₄P).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	O	P	0	0
			4	3	1		
5	A	1	Total	O	P	0	0
			4	3	1		

- Molecule 6 is NICKEL (II) ION (three-letter code: NI) (formula: Ni).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	1	Total Ni 1 1	0	0

- Molecule 7 is DELTA-2-ALBOMYCIN A1 (three-letter code: ALB) (formula: C₃₇H₅₇FeN₁₂O₁₈S).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	A	1	Total C Fe N O S 97 51 1 17 26 2	0	1


- Molecule 8 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
8	A	332	Total O 332 332	0	0

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: FERRIC HYDROXAMATE RECEPTOR



- Molecule 2: alpha-D-glucopyranose-(1-2)-alpha-D-glucopyranose-(1-3)-[alpha-D-galactopyranose-(1-6)]alpha-D-glucopyranose-(1-3)-[L-glycero-alpha-D-manno-heptopyranose-(1-7)]L-glycero-alpha-D-manno-heptopyranose-(1-3)-L-glycero-alpha-D-manno-heptopyranose-(1-5)-[3-deoxy-alpha-D-manno-oct-2-ulopyranosonic acid-(2-4)]3-deoxy-alpha-D-manno-oct-2-ulopyranosonic acid-(2-6)-2-amino-2,3-dideoxy-alpha-D-glucopyranose-(1-6)-2-amino-2-deoxy-alpha-D-glucopyranose

Chain B: 



4 Data and refinement statistics i

Property	Value	Source
Space group	P 61	Depositor
Cell constants a, b, c, α , β , γ	171.80Å 171.80Å 86.35Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	50.00 – 3.10 50.00 – 3.10	Depositor EDS
% Data completeness (in resolution range)	94.3 (50.00-3.10) 94.3 (50.00-3.10)	Depositor EDS
R_{merge}	0.15	Depositor
R_{sym}	0.14	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.16 (at 2.69Å)	Xtrriage
Refinement program	CNS 0.4	Depositor
R, R_{free}	0.222 , 0.283 0.224 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å ²)	71.3	Xtrriage
Anisotropy	0.327	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.28 , 60.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	0.028 for h,-h-k,-l	Xtrriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	6195	wwPDB-VP
Average B, all atoms (Å ²)	82.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.24% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GLC, PA1, GLA, PO4, KDO, FTT, GCN, GMH, ALB, NI, DPO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.92	0/5663	0.87	2/7696 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	A	179	ASP	CB-CG-OD2	-7.08	111.93	118.30
1	A	352	LEU	CA-CB-CG	5.02	126.85	115.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	599	TYR	Sidechain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5523	0	5223	516	0
2	B	134	0	102	11	0
3	A	84	0	126	8	0
4	A	16	0	0	1	0
5	A	8	0	0	2	0
6	A	1	0	0	0	0
7	A	97	0	36	13	0
8	A	332	0	0	67	0
All	All	6195	0	5487	542	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 48.

The worst 5 of 542 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:220:THR:HG23	1:A:230:THR:HG23	1.26	1.17
1:A:378:VAL:HG12	1:A:445:VAL:HG12	1.39	1.04
1:A:632:THR:HG23	1:A:642:THR:HG23	1.41	1.03
1:A:380:PHE:HD1	1:A:443:THR:HG23	1.22	1.02
7:A:1022[B]:ALB:H27	7:A:1022[B]:ALB:O11	1.54	1.02

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	705/725 (97%)	601 (85%)	77 (11%)	27 (4%)	2 15

5 of 27 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	24	PRO
1	A	224	ASP
1	A	412	HIS
1	A	696	ASP
1	A	25	ALA

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	587/601 (98%)	512 (87%)	75 (13%)	3 15

5 of 75 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	565	GLU
1	A	706	THR
1	A	600	THR
1	A	616	THR
1	A	273	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 18 such sidechains are listed below:

Mol	Chain	Res	Type
1	A	440	GLN
1	A	594	ASN
1	A	522	ASN
1	A	360	GLN
1	A	412	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

11 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	PA1	B	1	4,3,2	11,11,12	2.69	4 (36%)	14,15,17	2.69	4 (28%)
2	GMH	B	10	2	13,13,14	1.40	3 (23%)	17,18,20	0.62	0
2	KDO	B	11	2	15,15,16	1.08	1 (6%)	19,21,24	1.08	1 (5%)
2	GCN	B	2	3,2,5	10,10,11	2.34	5 (50%)	13,13,15	5.93	6 (46%)
2	KDO	B	3	2	15,15,16	1.04	1 (6%)	19,21,24	1.03	0
2	GMH	B	4	4,2	13,13,14	0.88	1 (7%)	17,18,20	0.94	1 (5%)
2	GMH	B	5	2,5	13,13,14	0.77	0	17,18,20	1.35	2 (11%)
2	GLC	B	6	2	11,11,12	0.44	0	15,15,17	1.03	1 (6%)
2	GLC	B	7	2	11,11,12	0.78	0	15,15,17	0.68	0
2	GLC	B	8	2	11,11,12	1.63	3 (27%)	15,15,17	1.19	2 (13%)
2	GLA	B	9	2	11,11,12	0.70	0	15,15,17	0.68	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	PA1	B	1	4,3,2	-	0/2/18/22	0/1/1/1
2	GMH	B	10	2	1/1/5/6	2/6/23/26	0/1/1/1
2	KDO	B	11	2	-	5/10/26/30	0/1/1/1
2	GCN	B	2	3,2,5	2/2/3/4	2/2/15/18	0/1/1/1
2	KDO	B	3	2	-	2/10/26/30	0/1/1/1
2	GMH	B	4	4,2	-	0/6/23/26	0/1/1/1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GMH	B	5	2,5	1/1/5/6	0/6/23/26	1/1/1/1
2	GLC	B	6	2	-	2/2/19/22	0/1/1/1
2	GLC	B	7	2	-	1/2/19/22	0/1/1/1
2	GLC	B	8	2	-	2/2/19/22	0/1/1/1
2	GLA	B	9	2	-	1/2/19/22	0/1/1/1

The worst 5 of 18 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	1	PA1	C1-C2	7.12	1.61	1.52
2	B	2	GCN	C3-C4	4.61	1.60	1.52
2	B	2	GCN	C1-C2	-3.29	1.48	1.52
2	B	2	GCN	C3-C2	-3.26	1.46	1.53
2	B	10	GMH	O5-C5	3.15	1.47	1.43

The worst 5 of 17 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	2	GCN	C3-C2-C1	19.67	125.80	109.82
2	B	1	PA1	C1-C2-N2	7.61	124.43	110.49
2	B	2	GCN	O4-C4-C5	5.30	121.25	110.01
2	B	5	GMH	C6-C5-C4	-3.90	107.57	114.03
2	B	1	PA1	C3-C4-C5	-3.80	105.83	110.77

All (4) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
2	B	2	GCN	C1
2	B	2	GCN	C2
2	B	5	GMH	C6
2	B	10	GMH	C6

5 of 17 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	11	KDO	O6-C6-C7-O7
2	B	11	KDO	O6-C6-C7-C8
2	B	8	GLC	O5-C5-C6-O6
2	B	8	GLC	C4-C5-C6-O6
2	B	2	GCN	O5-C5-C6-O6


All (1) ring outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	5	GMH	C1-C2-C3-C4-C5-O5

8 monomers are involved in 11 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	11	KDO	2	0
2	B	7	GLC	2	0
2	B	2	GCN	4	0
2	B	10	GMH	1	0
2	B	1	PA1	1	0
2	B	4	GMH	1	0
2	B	8	GLC	2	0
2	B	6	GLC	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.



5.6 Ligand geometry [i](#)

Of 13 ligands modelled in this entry, 1 is monoatomic - leaving 12 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	PO4	A	1014	2	0,3,4	-	-	0,3,6	-	-
3	FTT	A	1003	2	6,6,16	2.05	1 (16%)	6,7,17	2.12	1 (16%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	FTT	A	1002	2	15,15,16	2.12	2 (13%)	15,15,17	2.91	1 (6%)
5	PO4	A	1016	6,2	0,3,4	-	-	0,3,6	-	-
3	FTT	A	1004	3,2	15,15,16	1.70	2 (13%)	15,15,17	3.18	3 (20%)
7	ALB	A	1022[A]	-	72,75,75	2.32	17 (23%)	76,117,117	4.34	29 (38%)
7	ALB	A	1022[B]	-	72,75,75	2.43	15 (20%)	76,117,117	4.26	27 (35%)
3	FTT	A	1005	3	12,12,16	0.38	0	11,11,17	0.76	0
3	FTT	A	1006	3,2	16,16,16	1.32	1 (6%)	17,17,17	1.05	1 (5%)
4	DPO	A	1013	2	3,7,8	1.34	0	6,10,13	1.33	1 (16%)
3	FTT	A	1007	3	14,14,16	0.44	0	13,13,17	0.65	0
4	DPO	A	1015	2	3,7,8	1.18	0	6,10,13	0.88	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	FTT	A	1003	2	-	1/4/4/15	-
3	FTT	A	1002	2	-	1/14/14/15	-
3	FTT	A	1004	3,2	-	1/14/14/15	-
7	ALB	A	1022[A]	-	5/5/23/27	38/70/142/142	0/5/7/7
4	DPO	A	1013	2	-	0/2/5/6	-
3	FTT	A	1005	3	-	0/9/10/15	-
3	FTT	A	1006	3,2	1/1/2/2	1/15/15/15	-
7	ALB	A	1022[B]	-	7/7/23/27	33/70/142/142	0/5/7/7
3	FTT	A	1007	3	-	1/11/12/15	-
4	DPO	A	1015	2	-	0/2/5/6	-

The worst 5 of 38 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
7	A	1022[B]	ALB	C30-C29	9.60	1.66	1.54
7	A	1022[B]	ALB	C22-C24	9.28	1.77	1.52
7	A	1022[A]	ALB	C22-C24	8.23	1.74	1.52
7	A	1022[A]	ALB	C36-N8	-7.53	1.33	1.47
3	A	1002	FTT	C2-C1	7.43	1.69	1.49

The worst 5 of 63 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
7	A	1022[A]	ALB	C30-S-C31	18.68	115.53	94.41
7	A	1022[B]	ALB	C30-S-C31	17.70	114.42	94.41
7	A	1022[B]	ALB	C36-N8-C32	-15.50	90.13	117.31
7	A	1022[A]	ALB	C29-C30-S	-15.39	87.73	106.11
7	A	1022[B]	ALB	C29-C30-S	-12.70	90.94	106.11

5 of 13 chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
3	A	1006	FTT	C3
7	A	1022[A]	ALB	N3
7	A	1022[A]	ALB	N5
7	A	1022[A]	ALB	N4
7	A	1022[A]	ALB	C22

5 of 76 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
7	A	1022[A]	ALB	N2-C4-C8-C11
7	A	1022[A]	ALB	O4-C4-C8-C11
7	A	1022[A]	ALB	O4-C4-C8-N12
7	A	1022[A]	ALB	C8-C4-N2-C7
7	A	1022[A]	ALB	C14-C11-C8-N12


There are no ring outliers.


9 monomers are involved in 24 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	1014	PO4	2	0
3	A	1003	FTT	1	0
3	A	1004	FTT	4	0
7	A	1022[A]	ALB	5	0
7	A	1022[B]	ALB	8	0
3	A	1005	FTT	1	0
3	A	1006	FTT	6	0
3	A	1007	FTT	2	0
4	A	1015	DPO	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier.

Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	707/725 (97%)	-0.36	0 100 100	44, 80, 113, 117	0

There are no RSRZ outliers to report.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)


There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	GLC	B	8	11/12	0.56	0.16	116,116,117,118	0
2	GMH	B	10	13/14	0.75	0.15	115,116,117,118	0
2	GLC	B	7	11/12	0.80	0.10	111,114,116,117	0
2	GLA	B	9	11/12	0.82	0.10	110,112,113,113	0
2	GMH	B	5	13/14	0.88	0.08	89,98,111,116	0
2	GLC	B	6	11/12	0.89	0.08	93,101,110,113	0
2	KDO	B	11	15/16	0.90	0.07	80,84,88,90	0
2	GMH	B	4	13/14	0.94	0.07	72,78,83,87	0
2	PA1	B	1	11/12	0.94	0.06	53,64,78,81	0
2	KDO	B	3	15/16	0.95	0.07	67,74,78,84	0
2	GCN	B	2	10/11	0.97	0.05	46,54,63,71	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.



6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	FTT	A	1003	7/17	0.82	0.15	88,103,108,109	0
3	FTT	A	1002	16/17	0.88	0.17	80,91,96,96	0
5	PO4	A	1016	4/5	0.88	0.09	112,113,114,114	0
5	PO4	A	1014	4/5	0.91	0.11	76,77,78,84	0
4	DPO	A	1015	8/9	0.91	0.09	86,89,93,94	4
3	FTT	A	1006	17/17	0.92	0.14	67,91,100,100	0
3	FTT	A	1005	13/17	0.92	0.16	77,80,89,90	0
3	FTT	A	1007	15/17	0.93	0.16	79,99,107,110	0
3	FTT	A	1004	16/17	0.93	0.13	70,73,77,78	0
4	DPO	A	1013	8/9	0.94	0.08	84,86,90,92	4

Continued on next page...




Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
6	NI	A	1021	1/1	0.94	0.09	119,119,119,119	0
7	ALB	A	1022[A]	69/69	0.95	0.20	69,88,114,120	28
7	ALB	A	1022[B]	69/69	0.95	0.20	69,87,115,120	28

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around ALB A 1022 (A):

2mF_o-DF_c (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
 and green (positive)



Electron density around ALB A 1022 (B):

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.