



wwPDB EM Validation Summary Report ⓘ

Jun 11, 2025 – 12:40 AM JST

PDB ID : 8XLK / pdb_00008xlk
EMDB ID : EMD-38451
Title : Structure of native tri-heteromeric GluN1-GluN2A-GluN2B NMDA receptor
in rat cortex and hippocampus
Authors : Zhang, M.; Feng, J.; Li, Y.; Zhu, S.
Deposited on : 2023-12-26
Resolution : 4.20 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev118
Mogul : 1.8.5 (274361), CSD as541be (2020)
MolProbity : 4-5-2 with Phenix2.0rc1
buster-report : 1.1.7 (2018)
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
MapQ : 1.9.13
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.43.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

ELECTRON MICROSCOPY

The reported resolution of this entry is 4.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.










Metric	Whole archive (#Entries)	EM structures (#Entries)
Clashscore	210492	15764
Ramachandran outliers	207382	16835
Sidechain outliers	206894	16415

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	938	51% 34% 15%
1	C	938	54% 32% 15%
2	B	1464	36% 19% 46%
3	D	1482	34% 20% 47%
4	E	140	47% 34% 16%
4	I	140	49% 35% 16%
5	F	131	50% 34% 16%
5	J	131	50% 32% 16%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
6	G	227	
7	H	212	
8	K	248	
9	L	265	
10	M	3	
10	O	3	
11	N	2	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
13	7RC	B	1504	-	-	X	-

2 Entry composition [i](#)

There are 13 unique types of molecules in this entry. The entry contains 34118 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Glutamate receptor ionotropic, NMDA 1.

Mol	Chain	Residues	Atoms					AltConf	Trace
1	A	800	Total	C	N	O	S	0	0
			6322	4023	1099	1165	35		
1	C	800	Total	C	N	O	S	0	0
			6325	4025	1099	1165	36		

- Molecule 2 is a protein called Glutamate receptor ionotropic, NMDA 2A.

Mol	Chain	Residues	Atoms					AltConf	Trace
2	B	792	Total	C	N	O	S	0	0
			6279	4051	1025	1166	37		

- Molecule 3 is a protein called Glutamate receptor ionotropic, NMDA 2B.

Mol	Chain	Residues	Atoms					AltConf	Trace
3	D	792	Total	C	N	O	S	0	0
			6260	4026	1007	1184	43		

- Molecule 4 is a protein called Heavy Chain of GluN1 Fab, 4F11.

Mol	Chain	Residues	Atoms					AltConf	Trace
4	E	118	Total	C	N	O	S	0	0
			938	598	150	186	4		
4	I	118	Total	C	N	O	S	0	0
			938	598	150	186	4		

- Molecule 5 is a protein called Light Chain of GluN1 Fab, 4F11.

Mol	Chain	Residues	Atoms					AltConf	Trace
5	F	110	Total	C	N	O	S	0	0
			828	518	137	170	3		
5	J	110	Total	C	N	O	S	0	0
			822	512	137	170	3		

- Molecule 6 is a protein called Heavy Chain of GluN2A Fab, 28C.

Mol	Chain	Residues	Atoms					AltConf	Trace
6	G	227	Total	C	N	O	S	0	0
			1693	1061	294	334	4		

- Molecule 7 is a protein called Light Chain of GluN2A Fab, 28C.

Mol	Chain	Residues	Atoms					AltConf	Trace
7	H	212	Total	C	N	O	S	0	0
			1576	983	268	320	5		

- Molecule 8 is a protein called Heavy Chain of GluN2B Fab2.

Mol	Chain	Residues	Atoms					AltConf	Trace
8	K	116	Total	C	N	O	S	0	0
			899	569	152	173	5		

- Molecule 9 is a protein called Light Chain of GluN2B Fab2.

Mol	Chain	Residues	Atoms					AltConf	Trace
9	L	106	Total	C	N	O	S	0	0
			820	520	136	161	3		

- Molecule 10 is an oligosaccharide called alpha-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms					AltConf	Trace
10	M	3	Total	C	N	O		0	0
			39	22	2	15			
10	O	3	Total	C	N	O		0	0
			39	22	2	15			

- Molecule 11 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				AltConf	Trace
11	N	2	Total	C	N	O	0	0
			28	16	2	10		

- Molecule 12 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: $C_8H_{15}NO_6$).




Mol	Chain	Residues	Atoms				AltConf
12	A	1	Total	C	N	O	0
			14	8	1	5	
12	A	1	Total	C	N	O	0
			14	8	1	5	
12	A	1	Total	C	N	O	0
			14	8	1	5	
12	A	1	Total	C	N	O	0
			14	8	1	5	
12	A	1	Total	C	N	O	0
			14	8	1	5	
12	A	1	Total	C	N	O	0
			14	8	1	5	
12	B	1	Total	C	N	O	0
			14	8	1	5	
12	B	1	Total	C	N	O	0
			14	8	1	5	
12	B	1	Total	C	N	O	0
			14	8	1	5	

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				AltConf
12	C	1	Total	C	N	O	0
			14	8	1	5	
12	C	1	Total	C	N	O	0
			14	8	1	5	
12	C	1	Total	C	N	O	0
			14	8	1	5	
12	C	1	Total	C	N	O	0
			14	8	1	5	
12	C	1	Total	C	N	O	0
			14	8	1	5	
12	D	1	Total	C	N	O	0
			14	8	1	5	
12	D	1	Total	C	N	O	0
			14	8	1	5	
12	D	1	Total	C	N	O	0
			14	8	1	5	

- Molecule 13 is (2R)-4-(3-phosphonopropyl)piperazine-2-carboxylic acid (CCD ID: 7RC) (formula: C₈H₁₇N₂O₅P).



Mol	Chain	Residues	Atoms					AltConf
13	B	1	Total	C	N	O	P	0
			16	8	2	5	1	

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms					AltConf
			Total	C	N	O	P	
13	D	1	16	8	2	5	1	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Glutamate receptor ionotropic, NMDA 1



• Molecule 1: Glutamate receptor ionotropic, NMDA 1



• Molecule 2: Glutamate receptor ionotropic, NMDA 2A





ARG
ALA
PHE
ASN
GLY
SER
SER
ASN
GLY
HIS
VAL
TYR
GLU
LYS
LEU
SER
SER
ILE
GLU
SER
ASP
VAL

• Molecule 4: Heavy Chain of GluN1 Fab, 4F11

Chain E:



PRO MET VAL SER ILE VAL LEU TYR LEU LEU ALA ALA ALA HIS SER SER ILE GLU SER ASP VAL
D21 V22 G28 K33 L38 S39 L40 T41 C42 T43 Y47 S48 T49 T50 S51 D52 Y53 A54 W55 I58 R59 Q60 F61 P62 G63 N64 K65 L66 E67 W68 M69 G70 Y71

I72 S73 Y74 S75 G76 G79 N80 Y81 L84 I88 S89 I90 T91 R92 R93 T94 N97 Q98 F99 F100 L101 Q102 L103 N104 S105 D110 Y114 Y121 Y124 G132 V135 S138

• Molecule 4: Heavy Chain of GluN1 Fab, 4F11

Chain I:



PRO MET VAL SER ILE VAL LEU TYR LEU LEU ALA ALA ALA HIS SER SER ILE GLU SER ASP VAL
D21 V22 Q23 L24 E26 V32 S37 L38 C42 T43 V44 T45 D52 Y53 A54 I58 R59 L66 E67 W68 W69 G70 Y71 I72 S75 T78 G79 Y80

L84 K86 S86 R87 I88 T91 R92 D93 N97 Q98 F99 F100 L101 Q102 L103 N104 S105 V106 T107 T108 T111 A112 T113 Y114 R118 L119 L120 Y121 Y124 G125 M126 B127 Y128 Q131 T136 V137 S138

• Molecule 5: Light Chain of GluN1 Fab, 4F11

Chain F:



MET VAL SER ILE VAL LEU TYR VAL LEU LEU ALA ALA ALA HIS SER SER ILE GLU SER ASP VAL
I21 Q25 Q26 P27 L30 A31 V32 S33 Q36 R37 A38 T39 L40 V48 Y55 M56 N57 P58 T59 Q60 Q61 K62 Q65 K68 L69 L70 I71 Y72 A73 A74 S75 N76

R64 G67 T95 L96 H97 H99 P100 V101 E102 E103 Y109 Y110 C111 N115 E116 G123 G124 L127 K130

• Molecule 5: Light Chain of GluN1 Fab, 4F11

Chain J:



MET VAL SER ILE VAL LEU TYR VAL LEU LEU ALA ALA ALA HIS SER SER ILE GLU SER ASP VAL
I21 V22 L23 T24 Q25 A31 V32 S33 L34 G35 Q36 R37 A38 C42 K43 A44 S45 D49 N57 W58 Y59 Q60 Q61 K62 Q65 K68 L69 L70 I71 Y72 E78


G91 T92 I98 H99 P100 V101 A106 A107 T108 Y109 Y110 C111 Q112 Q113 G114 N115 P119 T120 F121 T125 K126 L127 K130

• Molecule 6: Heavy Chain of GluN2A Fab, 28C

Chain G:




G1 V2 T3 L4 K5 I11 L12 Q13 T17 L18 S23 F24 S25 A29 G33 V34 G37 W38 T39 R40 Q41 P42 K45 G46 L47 W49 H52 T53 A54 A55 Y61 K66 T70 D74 T75 Q79 K83 I84 A85 T89 T92 A93 T94




• Molecule 7: Light Chain of GluN2A Fab, 28C

Chain H: 67% 33%




• Molecule 8: Heavy Chain of GluN2B Fab2

Chain K: 6% 24% 23% 53%



• Molecule 9: Light Chain of GluN2B Fab2

Chain L: 6% 23% 17% 60%



- Molecule 10: alpha-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain M:  67% 33%

MAG1
MAG2
MAG3

- Molecule 10: alpha-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain O:  67% 33%

MAG1
MAG2
MAG3

- Molecule 11: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain N:  50% 50%

MAG1
MAG2

4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	45676	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	NONE	Depositor
Microscope	FEI POLARA 300	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	60	Depositor
Minimum defocus (nm)	1000	Depositor
Maximum defocus (nm)	2500	Depositor
Magnification	Not provided	
Image detector	DIRECT ELECTRON DE-10 (5k x 4k)	Depositor
Maximum map value	29.940	Depositor
Minimum map value	-20.122	Depositor
Average map value	0.000	Depositor
Map value standard deviation	1.000	Depositor
Recommended contour level	0.85	Depositor
Map size (\AA)	342.72, 342.72, 342.72	wwPDB
Map dimensions	320, 320, 320	wwPDB
Map angles ($^\circ$)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (\AA)	1.071, 1.071, 1.071	Depositor

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: 7RC, MAN, NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.13	0/6464	0.38	0/8757
1	C	0.13	0/6467	0.37	0/8760
2	B	0.13	0/6432	0.37	0/8736
3	D	0.14	0/6401	0.40	1/8674 (0.0%)
4	E	0.85	7/964 (0.7%)	1.00	8/1313 (0.6%)
4	I	0.15	0/964	0.43	0/1313
5	F	0.14	0/847	0.37	0/1152
5	J	0.54	3/840 (0.4%)	0.87	6/1143 (0.5%)
6	G	0.17	0/1727	0.49	0/2349
7	H	0.13	0/1607	0.37	0/2182
8	K	0.13	0/921	0.45	0/1247
9	L	0.15	0/840	0.47	0/1136
All	All	0.21	10/34474 (0.0%)	0.44	15/46762 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
4	E	0	1
9	L	0	1
All	All	0	2

The worst 5 of 10 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	E	52	ASP	CA-CB	-12.42	1.34	1.53
4	E	52	ASP	CA-C	-11.53	1.37	1.53
4	E	51	SER	CA-C	10.98	1.67	1.52
4	E	52	ASP	C-N	9.13	1.46	1.33

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	J	109	TYR	CA-C	8.18	1.63	1.52

The worst 5 of 15 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	E	52	ASP	N-CA-CB	-16.13	86.08	110.60
4	E	74	TYR	N-CA-C	-14.42	95.82	113.41
4	E	52	ASP	CA-C-O	-13.26	103.83	121.73
5	J	110	TYR	N-CA-C	9.96	125.62	109.59
5	J	107	ALA	O-C-N	-9.90	112.74	123.04

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
4	E	52	ASP	Mainchain
9	L	28	PRO	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	6322	0	6287	247	0
1	C	6325	0	6295	235	0
2	B	6279	0	6193	220	0
3	D	6260	0	6165	231	0
4	E	938	0	883	55	0
4	I	938	0	883	37	0
5	F	828	0	788	37	0
5	J	822	0	781	51	0
6	G	1693	0	1685	80	0
7	H	1576	0	1537	53	0
8	K	899	0	862	48	0
9	L	820	0	799	39	0
10	M	39	0	34	0	0
10	O	39	0	34	0	0
11	N	28	0	25	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
12	A	98	0	90	0	0
12	B	42	0	39	1	0
12	C	98	0	91	1	0
12	D	42	0	39	1	0
13	B	16	0	0	7	0
13	D	16	0	0	6	0
All	All	34118	0	33510	1244	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 18.

The worst 5 of 1244 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:D:762:TYR:OH	13:D:1504:7RC:C05	1.93	1.16
6:G:130:SER:HB2	6:G:153:ASN:O	1.55	1.05
1:C:410:TYR:HB2	1:C:456:TYR:O	1.59	1.02
1:C:350:ASN:HA	1:C:367:TYR:O	1.59	1.00
2:B:485:HIS:CG	13:B:1504:7RC:O05	2.18	0.97

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	796/938 (85%)	758 (95%)	37 (5%)	1 (0%)	48 82
1	C	796/938 (85%)	758 (95%)	36 (4%)	2 (0%)	37 71
2	B	788/1464 (54%)	743 (94%)	44 (6%)	1 (0%)	48 82
3	D	788/1482 (53%)	730 (93%)	57 (7%)	1 (0%)	48 82
4	E	116/140 (83%)	104 (90%)	9 (8%)	3 (3%)	4 28

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
4	I	116/140 (83%)	109 (94%)	7 (6%)	0	100	100
5	F	108/131 (82%)	96 (89%)	12 (11%)	0	100	100
5	J	108/131 (82%)	92 (85%)	14 (13%)	2 (2%)	6	35
6	G	225/227 (99%)	189 (84%)	35 (16%)	1 (0%)	30	67
7	H	210/212 (99%)	204 (97%)	6 (3%)	0	100	100
8	K	112/248 (45%)	106 (95%)	6 (5%)	0	100	100
9	L	104/265 (39%)	89 (86%)	15 (14%)	0	100	100
All	All	4267/6316 (68%)	3978 (93%)	278 (6%)	11 (0%)	38	71

5 of 11 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	659	VAL
1	C	545	GLU
4	E	53	TYR
6	G	225	PHE
1	A	811	GLU

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	689/806 (86%)	689 (100%)	0	100	100
1	C	690/806 (86%)	690 (100%)	0	100	100
2	B	695/1308 (53%)	695 (100%)	0	100	100
3	D	691/1299 (53%)	691 (100%)	0	100	100
4	E	104/119 (87%)	103 (99%)	1 (1%)	73	81
4	I	104/119 (87%)	104 (100%)	0	100	100
5	F	90/104 (86%)	90 (100%)	0	100	100
5	J	89/104 (86%)	88 (99%)	1 (1%)	70	79
6	G	181/181 (100%)	181 (100%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
7	H	169/169 (100%)	169 (100%)	0	100	100
8	K	95/205 (46%)	95 (100%)	0	100	100
9	L	91/221 (41%)	91 (100%)	0	100	100
All	All	3688/5441 (68%)	3686 (100%)	2 (0%)	92	94

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
4	E	51	SER
5	J	110	TYR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 35 such sidechains are listed below:

Mol	Chain	Res	Type
7	H	160	GLN
7	H	189	HIS
5	J	57	ASN
2	B	839	HIS
2	B	697	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

8 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
10	NAG	M	1	10,1	14,14,15	0.31	0	17,19,21	0.40	0
10	NAG	M	2	10	14,14,15	0.22	0	17,19,21	0.40	0
10	MAN	M	3	10	11,11,12	0.79	0	15,15,17	1.39	2 (13%)
11	NAG	N	1	2,11	14,14,15	0.66	1 (7%)	17,19,21	0.66	0
11	NAG	N	2	11	14,14,15	0.22	0	17,19,21	0.46	0
10	NAG	O	1	10,1	14,14,15	0.26	0	17,19,21	0.39	0
10	NAG	O	2	10	14,14,15	0.21	0	17,19,21	0.46	0
10	MAN	O	3	10	11,11,12	0.82	0	15,15,17	1.15	2 (13%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
10	NAG	M	1	10,1	-	2/6/23/26	0/1/1/1
10	NAG	M	2	10	-	2/6/23/26	0/1/1/1
10	MAN	M	3	10	-	2/2/19/22	1/1/1/1
11	NAG	N	1	2,11	-	2/6/23/26	0/1/1/1
11	NAG	N	2	11	-	2/6/23/26	0/1/1/1
10	NAG	O	1	10,1	-	1/6/23/26	0/1/1/1
10	NAG	O	2	10	-	2/6/23/26	0/1/1/1
10	MAN	O	3	10	-	2/2/19/22	1/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
11	N	1	NAG	O5-C1	-2.36	1.39	1.43

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
10	M	3	MAN	C1-O5-C5	4.16	117.83	112.19
10	O	3	MAN	C1-O5-C5	3.08	116.36	112.19
10	M	3	MAN	O2-C2-C3	-2.25	105.64	110.14
10	O	3	MAN	O2-C2-C3	-2.13	105.88	110.14

There are no chirality outliers.

5 of 15 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
10	O	2	NAG	O5-C5-C6-O6
10	M	2	NAG	O5-C5-C6-O6
10	M	1	NAG	O5-C5-C6-O6
10	O	3	MAN	O5-C5-C6-O6
10	O	2	NAG	C4-C5-C6-O6


All (2) ring outliers are listed below:


Mol	Chain	Res	Type	Atoms
10	O	3	MAN	C1-C2-C3-C4-C5-O5
10	M	3	MAN	C1-C2-C3-C4-C5-O5

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.







5.6 Ligand geometry [i](#)

22 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
12	NAG	C	1001	1	14,14,15	0.28	0	17,19,21	0.34	0
12	NAG	B	1501	2	14,14,15	0.25	0	17,19,21	0.43	0
12	NAG	A	1001	1	14,14,15	0.22	0	17,19,21	0.38	0
12	NAG	C	1007	1	14,14,15	0.19	0	17,19,21	0.37	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
12	NAG	A	1002	1	14,14,15	0.41	0	17,19,21	1.16	2 (11%)
12	NAG	C	1002	1	14,14,15	0.45	0	17,19,21	0.38	0
12	NAG	C	1006	1	14,14,15	0.21	0	17,19,21	0.38	0
12	NAG	A	1003	1	14,14,15	0.23	0	17,19,21	0.43	0
13	7RC	B	1504	-	15,16,16	2.00	1 (6%)	17,22,22	1.96	3 (17%)
12	NAG	B	1503	2	14,14,15	0.35	0	17,19,21	0.64	1 (5%)
12	NAG	A	1006	1	14,14,15	0.25	0	17,19,21	0.46	0
12	NAG	D	1502	3	14,14,15	0.25	0	17,19,21	0.40	0
12	NAG	B	1502	2	14,14,15	0.26	0	17,19,21	0.46	0
12	NAG	D	1503	3	14,14,15	0.41	0	17,19,21	1.16	2 (11%)
12	NAG	A	1004	1	14,14,15	0.23	0	17,19,21	0.41	0
13	7RC	D	1504	-	15,16,16	1.99	1 (6%)	17,22,22	1.79	5 (29%)
12	NAG	C	1005	1	14,14,15	0.27	0	17,19,21	0.36	0
12	NAG	A	1007	1	14,14,15	0.40	0	17,19,21	1.16	1 (5%)
12	NAG	C	1003	1	14,14,15	0.26	0	17,19,21	0.47	0
12	NAG	A	1005	1	14,14,15	0.24	0	17,19,21	0.42	0
12	NAG	C	1004	1	14,14,15	0.37	0	17,19,21	0.42	0
12	NAG	D	1501	3	14,14,15	0.25	0	17,19,21	0.42	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
12	NAG	C	1001	1	-	2/6/23/26	0/1/1/1
12	NAG	B	1501	2	-	2/6/23/26	0/1/1/1
12	NAG	A	1001	1	-	2/6/23/26	0/1/1/1
12	NAG	C	1007	1	-	4/6/23/26	0/1/1/1
12	NAG	A	1002	1	-	0/6/23/26	0/1/1/1
12	NAG	C	1002	1	-	0/6/23/26	0/1/1/1
12	NAG	C	1006	1	-	2/6/23/26	0/1/1/1
12	NAG	A	1003	1	-	2/6/23/26	0/1/1/1
13	7RC	B	1504	-	-	4/11/21/21	0/1/1/1
12	NAG	B	1503	2	-	0/6/23/26	0/1/1/1
12	NAG	A	1006	1	-	2/6/23/26	0/1/1/1
12	NAG	D	1502	3	-	2/6/23/26	0/1/1/1
12	NAG	B	1502	2	-	2/6/23/26	0/1/1/1
12	NAG	D	1503	3	-	0/6/23/26	0/1/1/1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
12	NAG	A	1004	1	-	2/6/23/26	0/1/1/1
13	7RC	D	1504	-	-	0/11/21/21	0/1/1/1
12	NAG	C	1005	1	-	2/6/23/26	0/1/1/1
12	NAG	A	1007	1	-	0/6/23/26	0/1/1/1
12	NAG	C	1003	1	-	2/6/23/26	0/1/1/1
12	NAG	A	1005	1	-	2/6/23/26	0/1/1/1
12	NAG	C	1004	1	-	1/6/23/26	0/1/1/1
12	NAG	D	1501	3	-	1/6/23/26	0/1/1/1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
13	B	1504	7RC	P01-C07	6.80	1.85	1.78
13	D	1504	7RC	P01-C07	6.75	1.85	1.78

The worst 5 of 14 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
13	B	1504	7RC	P01-C07-C06	-5.64	107.92	114.98
13	D	1504	7RC	P01-C07-C06	-4.21	109.71	114.98
13	D	1504	7RC	C03-N01-C02	2.84	113.86	109.52
13	B	1504	7RC	C04-N01-C02	-2.45	105.90	111.66
13	D	1504	7RC	C06-C04-N01	-2.37	107.88	113.84

There are no chirality outliers.

5 of 34 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
13	B	1504	7RC	C06-C07-P01-O03
13	B	1504	7RC	C06-C07-P01-O04
12	A	1006	NAG	O5-C5-C6-O6
12	A	1005	NAG	C4-C5-C6-O6
12	B	1502	NAG	O5-C5-C6-O6

There are no ring outliers.

5 monomers are involved in 16 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
12	C	1002	NAG	1	0
13	B	1504	7RC	7	0


Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Clashes	Symm-Clashes
12	B	1503	NAG	1	0
12	D	1502	NAG	1	0
13	D	1504	7RC	6	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-38451. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



X




Y




Z


6.1.2 Raw map



X



Y




Z


The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map



X Index: 160




Y Index: 160




Z Index: 160


6.2.2 Raw map



X Index: 160



Y Index: 160




Z Index: 160


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 154




Y Index: 176




Z Index: 151


6.3.2 Raw map



X Index: 155



Y Index: 176




Z Index: 150


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) [i](#)


6.4.1 Primary map



X




Y




Z


6.4.2 Raw map



X



Y




Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.

6.5 Orthogonal surface views [i](#)


6.5.1 Primary map



X




Y




Z

The images above show the 3D surface view of the map at the recommended contour level 0.85. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.


6.5.2 Raw map



X



Y



Z

These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.

6.6 Mask visualisation [i](#)

This section shows the 3D surface view of the primary map at 50% transparency overlaid with the specified mask at 0% transparency


A mask typically either:

- Encompasses the whole structure
- Separates out a domain, a functional unit, a monomer or an area of interest from a larger structure


6.6.1 emd_38451_msk_1.map [i](#)



X



Y




Z

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [i](#)



The volume at the recommended contour level is 2547 nm³; this corresponds to an approximate mass of 2300 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum ⓘ




*Reported resolution corresponds to spatial frequency of 0.238 \AA^{-1}

8 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC [i](#)



*Reported resolution corresponds to spatial frequency of 0.238 \AA^{-1}

8.2 Resolution estimates [i](#)

Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	4.20	-	-
Author-provided FSC curve	6.93	9.35	7.23
Unmasked-calculated*	6.85	9.22	7.19

*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from author-provided FSC intersecting FSC 0.143 CUT-OFF 6.93 differs from the reported value 4.2 by more than 10 %

The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 6.85 differs from the reported value 4.2 by more than 10 %

9 Map-model fit [i](#)


This section contains information regarding the fit between EMDB map EMD-38451 and PDB model 8XLK. Per-residue inclusion information can be found in section [3](#) on page [9](#).

9.1 Map-model overlay [i](#)




The images above show the 3D surface view of the map at the recommended contour level 0.85 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.85).

9.4 Atom inclusion [i](#)



At the recommended contour level, 99% of all backbone atoms, 98% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary ⓘ

The table lists the average atom inclusion at the recommended contour level (0.85) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	<div></div> 0.9790	<div></div> 0.2780
A	<div></div> 0.9950	<div></div> 0.3110
B	<div></div> 0.9930	<div></div> 0.3300
C	<div></div> 0.9930	<div></div> 0.3080
D	<div></div> 0.9920	<div></div> 0.2890
E	<div></div> 0.9760	<div></div> 0.1930
F	<div></div> 0.9990	<div></div> 0.1970
G	<div></div> 0.9530	<div></div> 0.2290
H	<div></div> 0.9860	<div></div> 0.2520
I	<div></div> 0.9880	<div></div> 0.1660
J	<div></div> 0.9980	<div></div> 0.2190
K	<div></div> 0.7550	<div></div> 0.0500
L	<div></div> 0.7600	<div></div> 0.0320
M	<div></div> 1.0000	<div></div> 0.3810
N	<div></div> 0.9290	<div></div> 0.3400
O	<div></div> 1.0000	<div></div> 0.3680

1.0

0.0

<0.0