



Full wwPDB X-ray Structure Validation Report ⓘ

Nov 3, 2024 – 01:14 AM EST

PDB ID : 1YQC
Title : Crystal Structure of Ureidoglycolate Hydrolase (AllA) from Escherichia coli O157:H7
Authors : Raymond, S.; Tocilj, A.; Matte, A.; Cygler, M.; Montreal-Kingston Bacterial Structural Genomics Initiative (BSGI)
Deposited on : 2005-02-01
Resolution : 1.71 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	1.20.1
EDS	:	3.0
Percentile statistics	:	20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4	:	9.0.003 (Gargrove)
Density-Fitness	:	1.0.11
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.39


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.71 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	5161 (1.70-1.70)
Clashscore	180529	5671 (1.70-1.70)
Ramachandran outliers	177936	5594 (1.70-1.70)
Sidechain outliers	177891	5594 (1.70-1.70)
RSRZ outliers	164620	5159 (1.70-1.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	170	<div> <div>11%</div> <div>68%</div> <div>24%</div> <div>6%</div> </div>
1	B	170	<div> <div>13%</div> <div>69%</div> <div>20%</div> <div>5%</div> <div>6%</div> </div>

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 2878 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Ureidoglycolate hydrolase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	160	Total	C	N	O	S	Se	0	0	0
			1277	813	220	240	2	2			
1	B	160	Total	C	N	O	S	Se	0	0	0
			1277	813	220	240	2	2			

There are 24 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	MSE	MET	modified residue	UNP P63486
A	84	MSE	MET	modified residue	UNP P63486
A	161	GLY	-	expression tag	UNP P63486
A	162	SER	-	expression tag	UNP P63486
A	163	HIS	-	expression tag	UNP P63486
A	164	HIS	-	expression tag	UNP P63486
A	165	HIS	-	expression tag	UNP P63486
A	166	HIS	-	expression tag	UNP P63486
A	167	HIS	-	expression tag	UNP P63486
A	168	HIS	-	expression tag	UNP P63486
A	169	HIS	-	expression tag	UNP P63486
A	170	HIS	-	expression tag	UNP P63486
B	1	MSE	MET	modified residue	UNP P63486
B	84	MSE	MET	modified residue	UNP P63486
B	161	GLY	-	expression tag	UNP P63486
B	162	SER	-	expression tag	UNP P63486
B	163	HIS	-	expression tag	UNP P63486
B	164	HIS	-	expression tag	UNP P63486
B	165	HIS	-	expression tag	UNP P63486
B	166	HIS	-	expression tag	UNP P63486
B	167	HIS	-	expression tag	UNP P63486
B	168	HIS	-	expression tag	UNP P63486
B	169	HIS	-	expression tag	UNP P63486
B	170	HIS	-	expression tag	UNP P63486

- Molecule 2 is GLYOXYLIC ACID (three-letter code: GLV) (formula: $C_2H_2O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			5	2	3		

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	161	Total	O	0	0
			161	161		
3	B	158	Total	O	0	0
			158	158		

4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	43.05Å 74.47Å 56.36Å 90.00° 103.89° 90.00°	Depositor
Resolution (Å)	36.44 – 1.71 36.44 – 1.71	Depositor EDS
% Data completeness (in resolution range)	95.5 (36.44-1.71) 95.4 (36.44-1.71)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.79 (at 1.71Å)	Xtriage
Refinement program	REFMAC refmac_5.2.0005	Depositor
R, R_{free}	0.226 , 0.269 0.225 , 0.268	Depositor DCC
R_{free} test set	1782 reflections (4.99%)	wwPDB-VP
Wilson B-factor (Å ²)	18.0	Xtriage
Anisotropy	0.060	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 41.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.55$, $\langle L^2 \rangle = 0.39$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	2878	wwPDB-VP
Average B, all atoms (Å ²)	22.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 45.69 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 1.2751e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GLV

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.90	1/1306 (0.1%)	1.01	5/1777 (0.3%)
1	B	0.91	2/1306 (0.2%)	1.00	6/1777 (0.3%)
All	All	0.90	3/2612 (0.1%)	1.00	11/3554 (0.3%)

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	118	TYR	CD1-CE1	5.65	1.47	1.39
1	A	16	TYR	CD1-CE1	5.42	1.47	1.39
1	B	16	TYR	CD1-CE1	5.34	1.47	1.39

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	140	ASP	CB-CG-OD2	7.60	125.14	118.30
1	B	37	ARG	NE-CZ-NH1	7.54	124.07	120.30
1	A	140	ASP	CB-CG-OD2	7.05	124.64	118.30
1	B	40	ASP	CB-CG-OD2	6.31	123.98	118.30
1	B	101	ASP	CB-CG-OD2	6.10	123.79	118.30
1	A	135	ASP	CB-CG-OD2	5.91	123.62	118.30
1	B	6	LEU	CA-CB-CG	5.79	128.62	115.30
1	A	148	ASP	CB-CG-OD2	5.45	123.20	118.30
1	A	97	ASP	CB-CG-OD2	5.24	123.02	118.30
1	A	26	ASP	CB-CG-OD2	5.16	122.95	118.30
1	B	98	ASP	CB-CG-OD2	5.13	122.91	118.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1277	0	1236	46	0
1	B	1277	0	1236	44	0
2	A	5	0	1	0	0
3	A	161	0	0	14	0
3	B	158	0	0	11	0
All	All	2878	0	2473	75	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 15.

All (75) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:142:GLY:HA2	1:B:37:ARG:NH2	1.47	1.25
1:A:37:ARG:NH2	1:B:142:GLY:HA2	1.52	1.25
1:B:47:LEU:HD21	1:B:120:ARG:NH2	1.64	1.11
1:A:142:GLY:CA	1:B:37:ARG:HH21	1.64	1.09
1:A:137:LEU:HD23	1:B:137:LEU:HD23	1.33	1.06
1:B:71:LEU:HD23	3:B:289:HOH:O	1.58	1.03
1:A:142:GLY:HA2	1:B:37:ARG:HH21	0.96	1.02
1:A:37:ARG:HH21	1:B:142:GLY:HA2	0.85	1.00
1:A:37:ARG:HH21	1:B:142:GLY:CA	1.73	0.99
1:A:1:MSE:HA	3:A:1109:HOH:O	1.66	0.96
1:B:47:LEU:HD21	1:B:120:ARG:HH21	1.18	0.94
1:A:99:LYS:HG2	3:A:1120:HOH:O	1.78	0.83
1:A:1:MSE:CA	3:A:1109:HOH:O	2.24	0.81
1:B:10:GLN:HE22	1:B:19:VAL:H	1.25	0.81
1:A:137:LEU:CD2	1:B:137:LEU:HD23	2.11	0.80
1:A:142:GLY:CA	1:B:37:ARG:NH2	2.30	0.78
1:A:82:ILE:HD13	1:B:137:LEU:HD21	1.66	0.77
1:A:10:GLN:HE22	1:A:19:VAL:H	1.33	0.77
1:A:2:LYS:HE3	1:A:156:GLU:HB3	1.67	0.74
1:A:49:GLN:O	3:A:1156:HOH:O	2.08	0.72
1:B:47:LEU:HD22	1:B:47:LEU:H	1.54	0.71
1:A:137:LEU:HD23	1:B:137:LEU:CD2	2.15	0.71

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:73:ARG:NH2	1:A:97:ASP:O	2.23	0.70
1:B:29:HIS:HE1	3:B:233:HOH:O	1.76	0.69
1:B:10:GLN:NE2	1:B:19:VAL:H	1.90	0.68
1:A:137:LEU:HD21	1:B:82:ILE:HD13	1.76	0.68
1:B:125:HIS:HD2	3:B:289:HOH:O	1.76	0.68
1:A:11:GLU:HG3	3:A:1127:HOH:O	1.96	0.66
1:A:10:GLN:NE2	1:A:19:VAL:H	1.93	0.66
1:B:64:LEU:HA	1:B:65:PRO:C	2.18	0.64
1:B:11:GLU:H	1:B:11:GLU:CD	2.01	0.62
1:B:71:LEU:CD2	3:B:289:HOH:O	2.29	0.61
1:B:48:GLU:OE2	1:B:141:ARG:NH2	2.34	0.61
1:B:79:GLN:NE2	1:B:124:HIS:HE1	1.98	0.60
1:A:64:LEU:HD13	3:A:1147:HOH:O	2.02	0.59
1:A:37:ARG:NH2	1:B:142:GLY:CA	2.45	0.57
1:A:29:HIS:HE1	3:A:1091:HOH:O	1.87	0.56
1:B:125:HIS:CD2	3:B:289:HOH:O	2.54	0.56
1:A:98:ASP:HA	3:A:1104:HOH:O	2.06	0.55
1:B:48:GLU:CD	1:B:141:ARG:HH21	2.10	0.55
1:B:48:GLU:CD	1:B:141:ARG:NH2	2.60	0.54
1:B:79:GLN:HE21	1:B:124:HIS:CE1	2.26	0.54
1:A:52:THR:HB	3:A:1156:HOH:O	2.07	0.54
1:A:64:LEU:HA	1:A:65:PRO:C	2.30	0.52
1:A:150:GLU:CD	3:A:1120:HOH:O	2.47	0.51
1:B:49:GLN:HE21	1:B:49:GLN:HA	1.76	0.51
1:B:79:GLN:NE2	1:B:124:HIS:CE1	2.78	0.49
1:B:47:LEU:HD22	1:B:47:LEU:N	2.26	0.49
1:B:106:ARG:NE	3:B:271:HOH:O	2.46	0.48
1:A:74:HIS:HD2	3:A:1100:HOH:O	1.95	0.48
1:A:73:ARG:HD3	1:A:148:ASP:OD1	2.15	0.47
1:A:142:GLY:HA3	1:B:37:ARG:HH21	1.68	0.47
1:B:92:VAL:O	1:B:124:HIS:HA	2.15	0.47
1:B:98:ASP:HB2	3:B:290:HOH:O	2.15	0.46
1:A:73:ARG:HD2	1:A:123:TRP:CH2	2.49	0.46
1:B:141:ARG:NH1	1:B:145:ASP:OD2	2.48	0.46
1:A:23:GLN:NE2	1:A:111:ASN:HD21	2.14	0.46
1:A:23:GLN:HG3	3:A:1110:HOH:O	2.15	0.45
1:A:133:VAL:HG23	3:A:1035:HOH:O	2.16	0.45
1:B:45:GLU:HB2	1:B:78:THR:HG23	1.97	0.45
1:A:52:THR:HG21	3:B:273:HOH:O	2.15	0.45
1:B:104:THR:HA	3:B:310:HOH:O	2.18	0.43
1:B:80:ALA:HB2	1:B:117:ASN:HD22	1.84	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:34:LEU:HD12	1:A:34:LEU:HA	1.75	0.43
1:A:102:LEU:HD22	1:A:102:LEU:H	1.84	0.43
1:A:45:GLU:HG2	3:B:295:HOH:O	2.19	0.43
1:A:57:ASN:HD21	1:B:37:ARG:HH11	1.65	0.42
1:A:8:LEU:HD22	1:A:110:THR:HG22	2.02	0.42
1:B:23:GLN:HB3	3:B:318:HOH:O	2.18	0.42
1:A:126:PRO:HG2	3:A:1155:HOH:O	2.19	0.42
1:A:64:LEU:HB2	1:A:65:PRO:HA	2.01	0.41
1:A:51:ARG:O	1:A:141:ARG:HA	2.20	0.41
1:A:92:VAL:O	1:A:124:HIS:HA	2.20	0.41
1:B:102:LEU:HD13	1:B:102:LEU:HA	1.68	0.40
1:A:15:ALA:O	1:A:95:LEU:HD21	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	158/170 (93%)	152 (96%)	6 (4%)	0	100	100
1	B	158/170 (93%)	151 (96%)	7 (4%)	0	100	100
All	All	316/340 (93%)	303 (96%)	13 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	139/148 (94%)	135 (97%)	4 (3%)	37	20
1	B	139/148 (94%)	127 (91%)	12 (9%)	8	2
All	All	278/296 (94%)	262 (94%)	16 (6%)	17	5

All (16) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1	MSE
1	A	34	LEU
1	A	145	ASP
1	A	151	SER
1	B	3	LEU
1	B	34	LEU
1	B	47	LEU
1	B	49	GLN
1	B	53	LEU
1	B	64	LEU
1	B	66	LEU
1	B	67	THR
1	B	73	ARG
1	B	102	LEU
1	B	120	ARG
1	B	141	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (14) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	10	GLN
1	A	29	HIS
1	A	57	ASN
1	A	60	GLN
1	A	74	HIS
1	A	79	GLN
1	A	117	ASN
1	B	10	GLN
1	B	23	GLN
1	B	29	HIS
1	B	49	GLN
1	B	79	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	117	ASN
1	B	124	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	GLV	A	1001	-	4,4,4	1.66	1 (25%)	3,4,4	1.32	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GLV	A	1001	-	-	0/0/2/2	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1001	GLV	O2-C2	2.34	1.28	1.23

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	158/170 (92%)	0.66	19 (12%) 10 9	10, 19, 35, 38	0
1	B	158/170 (92%)	0.79	22 (13%) 7 7	11, 19, 36, 41	0
All	All	316/340 (92%)	0.72	41 (12%) 9 8	10, 19, 36, 41	0

All (41) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	49	GLN	5.0
1	A	146	ASN	4.3
1	B	146	ASN	4.3
1	B	47	LEU	4.0
1	B	33	GLY	3.9
1	A	64	LEU	3.8
1	B	50	ASP	3.6
1	B	143	GLY	3.5
1	A	63	ASN	3.5
1	B	160	ALA	3.2
1	B	64	LEU	3.0
1	B	145	ASP	2.8
1	B	34	LEU	2.8
1	A	145	ASP	2.7
1	A	151	SER	2.7
1	B	102	LEU	2.7
1	A	6	LEU	2.7
1	A	34	LEU	2.6
1	B	65	PRO	2.6
1	A	160	ALA	2.5
1	A	33	GLY	2.4
1	A	149	VAL	2.4
1	B	158	CYS	2.4
1	B	37	ARG	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	98	ASP	2.3
1	A	96	GLY	2.3
1	B	120	ARG	2.3
1	A	4	GLN	2.2
1	A	62	ALA	2.2
1	A	31	ASN	2.2
1	A	65	PRO	2.2
1	A	66	LEU	2.2
1	B	144	SER	2.1
1	A	102	LEU	2.1
1	B	149	VAL	2.1
1	A	158	CYS	2.1
1	B	95	LEU	2.1
1	B	142	GLY	2.1
1	B	66	LEU	2.1
1	B	3	LEU	2.0
1	B	151	SER	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	GLV	A	1001	5/5	0.74	0.17	31,35,35,42	0

6.5 Other polymers [i](#)

There are no such residues in this entry.